Author Index

Ahnoff, M., see Hansson, G. 666(1994)505

Allenmark, S.G. and Andersson, S.
Proteins and peptides as chiral selectors in liquid chromatography (Review) 666(1994)167

Andersson, S., see Allenmark, S.G. 666(1994)167

Angelo, H.R., see Kristensen, K. 666(1994)283

Anton, K., Eppinger, J., Frederiksen, L., Francotte, E., Berger, T.A. and Wilson, W.H.
Chiral separations by packed-column super- and subcritical fluid chromatography 666(1994)395

Aoki, F., see Ôi, N. 666(1994)457

Araki, T., see Hosoya, K. 666(1994)449

Armstrong, D.W. and Zukowski, J.
Direct enantiomeric resolution of monoterpene hydrocarbons via reversed-phase high-performance liquid chromatography with an α-cyclodextrin bonded stationary phase 666(1994)445

Armstrong, D.W., see Pawlowska, M. 666(1994)485

Armstrong, D.W., see Tang, Y. 666(1994)147

Asakawa, N., see Ishihama, Y. 666(1994)193

Bargmann-Leyder, N., Truffert, J.-C., Tambuté, A. and Caude, M.
Evaluation of Pirkle-type chiral stationary phases by liquid and supercritical fluid chromatography. Influence of the spacer length and the steric hindrance in the vicinity of the stereogenic centre 666(1994)27

Belder, D. and Schomburg, G.
Chiral separations of basic and acidic compounds in modified capillaries using cyclodextrin-modified capillary zone electrophoresis 666(1994)351

Bereuter, T., see Kuhn, R. 666(1994)367

Berger, T.A., see Anton, K. 666(1994)395

Bertucci, C., Salvadori, P. and Lopes Guimaraes, L.F.
Determination of optical purity by high-performance liquid chromatography upon non-chiral stationary phases with dual circular dichroism/absorption detection 666(1994)535

Bicchi, C., D'Amato, A., Manzin, V., Galli, A. and Galli, M.
Cyclodextrin derivatives in the gas chromatographic separation of racemic mixtures of volatile compounds. VII. The use of 2,6-di-O-methyl-3 O-pentyl-β-cyclodextrin diluted in phases with different polarity in the separation of racemates in complex mixtures 666(1994)137

Blaschke, E., see Egginger, G. 666(1994)275

Blaschke, G., see Knoche, B. 666(1994)235

Blemmer, T., see Kristensen, K. 666(1994)283

Bossù, E., see Quaglia, M.G. 666(1994)289

Brückner, H., Haasmann, S., Langer, M., Westhauser, T., Wittner, R. and Godel, H.
Liquid chromatographic determination of D- and L-amino acids by derivatization with *o*-phthaldialdehyde and chiral thiols. Applications with reference to biosciences 666(1994)259

Buda, W.M., see Jaques, K. 666(1994)131

Cabrera, K. and Lubda, D.
Influence of temperature on chiral high-performance liquid chromatographic separations of oxazepam and Prominal on chemically bonded β-cyclodextrin as stationary phase 666(1994)433

Caude, M., see Bargmann-Leyder, N. 666(1994)27

Corradini, R., see Galli, B. 666(1994)77

Cvak, L., see Sinibaldi, M. 666(1994)471

D'Amato, A., see Bicchi, C. 666(1994)137

Davankov, V.A.
Chiral selectors with chelating properties in liquid chromatography: fundamental reflections and selective review of recent developments (Review) 666(1994)55

Davankov, V., see Kurganov, A. 666(1994)99

Desideri, N., see Quaglia, M.G. 666(1994)289

Dietrich, A., see Faber, B. 666(1994)161

Dingenen, J. and Kinkel, J.N.
Preparative chromatographic resolution of racemates on chiral stationary phases on laboratory and production scales by closed-loop recycling chromatography 666(1994)627

Dossena, A., see Galli, B. 666(1994)77

Dukesherer, D.R., see Hamper, B.C. 666(1994)479

Edholm, L.-E., see Pálmarsdóttir, S. 666(1994)337

Egginger, G., Blaschke, E., Lindner, W. and Olsson, A.-M.
Stereoselective high-performance liquid chromatographic assay of (±)-delmopinol in plasma using solid-phase extraction, a chiral derivatizing agent and electrochemical detection 666(1994)275

Eisenbeiss, F., see Kurganov, A. 666(1994)99

Eppinger, J., see Anton, K. 666(1994)395

Erni, F., see Kuhn, R. 666(1994)367

Erni, F., see Werner, A. 666(1994)375

Faber, B., Dietrich, A. and Mosandl, A.
Chiral compounds of essential oils. XV. Stereodifferentiation of characteristic compounds of *Mentha* species by multi-dimensional gas chromatography 666(1994)161

Farina, A., see Quaglia, M.G. 666(1994)289

Farkas, G., see Irgens, L.H. 666(1994)603

Flieger, M., see Sinibaldi, M. 666(1994)471

Floersheim, Ph., see Küsters, E. 666(1994)421

Francotte, E.
Contribution of preparative chromatographic resolution to the investigation of chiral phenomena (Review) 666(1994)565

Francotte, E., see Anton, K. 666(1994)395

Fréchet, J.M.J., see Hosoya, K. 666(1994)449

Frederiksen, L., see Anton, K. 666(1994)395

Galli, A., see Bicchi, C. 666(1994)137

Galli, B., Gasparrini, F., Misiti, D., Villani, C., Corradini, R., Dossena, A. and Marchelli, R.
Enantiomeric separation of dansyl- and dabsylamino acids by ligand-exchange chromatography with (*S*)- and (*R*)-phenylalaninamide-modified silica gel 666(1994)77

Galli, M., see Bicchi, C. 666(1994)137
Gasparrini, F., see Galli, B. 666(1994)77
Gil-Av, E. and Schurig, V.
Resolution of non-racemic mixtures in achiral chromatographic systems: a model for the enantioselective effects observed 666(1994)519
Godel, H., see Brückner, H. 666(1994)259
Graf, R., see Koppenhoefer, B. 666(1994)557
Grosenick, H., see Schurig, V. 666(1994)617
Gübitz, G., Mihellyes, S., Kobinger, G. and Wutte, A.
New chemically bonded chiral ligand-exchange chromatographic stationary phases 666(1994)91
Haas, P., see Kuhn, R. 666(1994)367
Haasmann, S., see Brückner, H. 666(1994)259
Hacksell, U., see Luthman, K. 666(1994)527
Haginaka, J., Murashima, T. and Seyama, C.
Separation of enantiomers on a lysozyme-bonded silica column 666(1994)203
Hamper, B.C., Dukesherer, D.R. and Moedritzer, K.
Analytical and preparative separation of the enantiomers of pyrazole phenyl ether herbicides on three chiral stationary phases 666(1994)479
Hansson, G. and Ahnoff, M.
Chromatographic separation of amide diastereomers: correlation with molecular descriptors 666(1994)505
Hardt, I. and König, W.A.
Preparative enantiomer separation with modified cyclodextrins as chiral stationary phases 666(1994)611
Hermansson, I., see Hermansson, J. 666(1994)181
Hermansson, J. and Hermansson, I.
Dynamic modification of the chiral bonding properties of a CHIRAL-AGP column by organic and inorganic additives. Separation of enantiomers of anti-inflammatory drugs 666(1994)181
Hobo, T., see Sato, K. 666(1994)463
Holzschuh, H., see Koppenhoefer, B. 666(1994)557
Hosoya, K., Yoshizako, K., Tanaka, N., Kimata, K., Araki, T. and Fréchet, J.M.J.
One-pot preparation method for a uniform-sized polymer-based chiral stationary phase for high-performance liquid chromatography with polymethacrylamide as a chiral selector 666(1994)449
Irgens, L.H., Farkas, G. and Vigh, G.
Displacement chromatography on cyclodextrin silicas. V. Separation of the enantiomers of 5,10-dideazatetrahydrofolic acid 666(1994)603
Ishihama, Y., Oda, Y., Asakawa, N., Yoshida, Y. and Sato, T.
Optical resolution by electrokinetic chromatography using ovomucoid as a pseudo-stationary phase 666(1994)193
Jaques, K., Buda, W.M., Venema, A. and Sandra, P.
Enantioselective separations by capillary gas chromatography on derivatized cyclodextrins. III. Separation of some racemic 2,2-dialkyl-4-alkoxycarbonyl-1,3-dioxolane derivatives on 2,3-di-O-acetyl-6-O-*tert*.-butyldimethylsilyl-β- and -γ-cyclodextrins 666(1994)131
Jensen, A.V., see Luthman, K. 666(1994)527
Kaida, Y., see Okamoto, Y. 666(1994)403
Karlsson, A., see Luthman, K. 666(1994)527
Kiechle, P., see Werner, A. 666(1994)375
Kimata, K., see Hosoya, K. 666(1994)449
Kinkel, J., see Kurganov, A. 666(1994)99
Kinkel, J.N., see Dingenen, J. 666(1994)627
Kirkland, K.M. and McCombs, D.A.
Changes in chiral selectivity with temperature for an ovomucoid protein-based column 666(1994)211
Kitahara, H., see Ôi, N. 666(1994)457
Knoche, B. and Blaschke, G.
Investigations on the *in vitro* racemization of thalidomide by high-performance liquid chromatography 666(1994)235
Kobinger, G., see Gübitz, G. 666(1994)91
König, W.A., see Hardt, I. 666(1994)611
Koppenhoefer, B., Graf, R., Holzschuh, H., Nothdurft, A., Trettin, U., Piras, P. and Roussel, C.
CHIRBASE, a molecular database for the separation of enantiomers by chromatography 666(1994)557
Krause, H., see Winkler, F.J. 666(1994)549
Kristensen, K., Angelo, H.R. and Blemmer, T.
Enantioselective high-performance liquid chromatographic method for the determination of methadone in serum using an AGP and a CN column as chiral and analytical column, respectively 666(1994)283
Kuhn, R., Steinmetz, C., Bereuter, T., Haas, P. and Erni, F.
Enantiomeric separations in capillary zone electrophoresis using a chiral crown ether 666(1994)367
Kurganov, A., Davankov, V., Unger, K., Eisenbeiss, F. and Kinkel, J.
Unusual peak distortion in ligand-exchange chromatography of enantiomers under overloaded conditions 666(1994)99
Küsters, E., Loux, V., Schmid, E. and Floersheim, Ph.
Enantiomeric separation of chiral sulphoxides. Screening of cellulose-based sorbents with particular reference to cellulose tribenzoate 666(1994)421
Langer, M., see Brückner, H. 666(1994)259
Li, S. and Lloyd, D.K.
Packed-capillary electrochromatographic separation of the enantiomers of neutral and anionic compounds using β-cyclodextrin as a chiral selector. Effect of operating parameters and comparison with free-solution capillary electrophoresis 666(1994)321
Lindner, W.
Preface 666(1994)1
Lindner, W., see Egginger, G. 666(1994)275
Lindner, W., see Uray, G. 666(1994)41
Lipkowitz, K.B.
Theoretical studies of brush-type chiral stationary phases (Review) 666(1994)493
Lloyd, D.K., see Li, S. 666(1994)321
Lopes Guimaraes, L.F., see Bertucci, C. 666(1994)535
Loux, V., see Küsters, E. 666(1994)421
Lu, X.-L. and Yang, S.K.
Enantiomer resolution of camazepam and its derivatives and enantioselective metabolism of camazepam by human liver microsomes 666(1994)249
Lubda, D., see Cabrera, K. 666(1994)433

Luthman, K., Jensen, A.V., Hacksell, U., Karlsson, A. and Pettersson, C.
Chiral discrimination of complexes between benzyloxycarbonylglycyl-L-proline and 4-hydroxy-2-dipropylaminoindan in ion-pair chromatography. A molecular mechanics study 666(1994)527
Maier, N.M., see Uray, G. 666(1994)41
Manzin, V., see Bicchi, C. 666(1994)137
Marchelli, R., see Galli, B. 666(1994)77
Markides, K.E., see Petersson, P. 666(1994)381
McCombs, D.A., see Kirkland, K.M. 666(1994)211
Medina, R., see Winkler, F.J. 666(1994)549
Messina, A., see Sinibaldi, M. 666(1994)471
Mihellyes, S., see Gübitz, G. 666(1994)91
Misiti, D., see Galli, B. 666(1994)77
Miwa, T., see Takeuchi, T. 666(1994)439
Moedritzer, K., see Hamper, B.C. 666(1994)479
Morrone, L.A., see Quaglia, M.G. 666(1994)289
Mosandl, A., see Faber, B. 666(1994)161
Murashima, T., see Haginaka, J. 666(1994)203
Muschalek, V., see Zimmer, D. 666(1994)241
Nakano, H., see Sato, K. 666(1994)463
Nassauer, T., see Werner, A. 666(1994)375
Nishi, H., see Terabe, S. 666(1994)295
Nothdurft, A., see Koppenhoefer, B. 666(1994)557
Oda, Y., see Ishihama, Y. 666(1994)193
Ôi, N., Kitahara, H. and Aoki, F.
Enantiomer separation by high-performance liquid chromatography with (*R*,*R*)-tartaric acid mono-amide derivatives as bifunctional chiral selectors 666(1994)457
Okamoto, Y. and Kaida, Y.
Resolution by high-performance liquid chromatography using polysaccharide carbamates and benzoates as chiral stationary phases (Review) 666(1994)403
Olsson, A.-M., see Egginger, G. 666(1994)275
Otsuka, K., see Terabe, S. 666(1994)295
Pálmarsdóttir, S. and Edholm, L.-E.
Capillary zone electrophoresis for separation of drug enantiomers using cyclodextrins as chiral selectors. Influence of experimental parameters on separation 666(1994)337
Pawlowska, M., Zukowski, J. and Armstrong, D.W.
Sensitive enantiomeric separation of aliphatic and aromatic amines using aromatic anhydrides as non-chiral derivatizing agents 666(1994)485
Pettersson, C., see Luthman, K. 666(1994)527
Petersson, P. and Markides, K.E.
Chiral separations performed by supercritical fluid chromatography (Review) 666(1994)381
Pichini, A., see Sinibaldi, M. 666(1994)471
Pini, D., see Uccello-Barretta, G. 666(1994)541
Piras, P., see Koppenhoefer, B. 666(1994)557
Quaglia, M.G., Desideri, N., Bossù, E., Farina, A. and Morrone, L.A.
Chromatographic resolution and pharmacological investigation of ICI 118551, a new β_2-blocker 666(1994)289
Rosini, C., see Uccello-Barretta, G. 666(1994)541
Roussel, C., see Koppenhoefer, B. 666(1994)557
Salvadori, P., see Bertucci, C. 666(1994)535
Salvadori, P., see Uccello-Barretta, G. 666(1994)541
Sandra, P., see Jaques, K. 666(1994)131
Sato, K., Nakano, H. and Hobo, T.
Effect of sample derivatization in liquid chromatographic separations of amine and amino acid enantiomers using diamide-type stationary phases 666(1994)463
Sato, T., see Ishihama, Y. 666(1994)193
Schmid, E., see Küsters, E. 666(1994)421
Schomburg, G., see Belder, D. 666(1994)351
Schurig, V.
Enantiomer separation by gas chromatography on chiral stationary phases (Review) 666(1994)111
Schurig, V. and Grosenick, H.
Preparative enantiomer separation of enflurane and isoflurane by inclusion gas chromatography 666(1994)617
Schurig, V., see Gil-Av, E. 666(1994)519
Seyama, C., see Haginaka, J. 666(1994)203
Sinibaldi, M., Flieger, M., Cvak, L., Messina, A. and Pichini, A.
Direct resolution of optically active isomers on chiral packings containing ergoline skeletons. II. Enantioseparation of carboxylic acids 666(1994)471
Steinmetz, C., see Kuhn, R. 666(1994)367
Takeuchi, T. and Miwa, T.
Separation of ferrocenylethanol enantiomers by microcolumn liquid chromatography with β-cyclodextrin as mobile phase additive 666(1994)439
Tambuté, A., see Bargmann-Leyder, N. 666(1994)27
Tanaka, N., see Hosoya, K. 666(1994)449
Tang, Y., Zhou, Y. and Armstrong, D.W.
Examination of the enantioselectivity of wall-immobilized cyclodextrin copolymers in capillary gas chromatography 666(1994)147
Terabe, S., Otsuka, K. and Nishi, H.
Separation of enantiomers by capillary electrophoretic techniques (Review) 666(1994)295
Trettin, U., see Koppenhoefer, B. 666(1994)557
Truffert, J.-C., see Bargmann-Leyder, N. 666(1994)27
Uccello-Barretta, G., Pini, D., Rosini, C. and Salvadori, P.
Stereochemical features of the stereoselective interaction between (*R*)- or (*S*)-7,7′-bis(1-propen-3-oxy)-2,2′-dihydroxy-1,1′-binaphthyl and quinine 666(1994)541
Unger, K., see Kurganov, A. 666(1994)99
Uray, G., Maier, N.M. and Lindner, W.
Diphenylethanediamine derivatives as chiral selectors. III . Comparison of four new diastereomeric chiral stationary phases prepared by addition of mono-3,5-dinitrobenzoyldiphenylethanediamine derivatives to optically pure epoxy silica 666(1994)41
Venema, A., see Jaques, K. 666(1994)131
Vigh, G., see Irgens, L.H. 666(1994)603
Villani, C., see Galli, B. 666(1994)77
Wainer, I.W.
Enantioselective high-performance liquid affinity chromatography as a probe of ligand–biopolymer interactions: an overview of a different use for high-performance liquid chromatographic chiral stationary phases 666(1994)221

Welch, C.J.
Evolution of chiral stationary phase design in the Pirkle laboratories (Review) 666(1994)3
Werner, A., Nassauer, T., Kiechle, P. and Erni, F.
Chiral separation by capillary zone electrophoresis of an optically active drug and amino acids by host–guest complexation with cyclodextrins 666(1994)375
Westhauser, T., see Brückner, H. 666(1994)259
Wilson, W.H., see Anton, K. 666(1994)395
Winkler, F.J., Medina, R., Winkler, J. and Krause, H.
Mass spectral and semi-empirical studies on chiral discrimination in gaseous aggregation products of protonated dialkyl tartrates 666(1994)549
Winkler, J., see Winkler, F.J. 666(1994)549
Wittner, R., see Brückner, H. 666(1994)259
Wutte, A., see Gübitz, G. 666(1994)91
Yang, S.K., see Lu, X.-L. 666(1994)249
Yoshida, Y., see Ishihama, Y. 666(1994)193
Yoshizako, K., see Hosoya, K. 666(1994)449
Zhou, Y., see Tang, Y. 666(1994)147
Zimmer, D. and Muschalek, V.
Enantioselective assay for the determination of nisoldipine in dog, rat and mouse plasma by chiral microbore high-performance liquid chromatography combined with gas chromatography–mass spectrometry 666(1994)241
Zukowski, J., see Armstrong, D.W. 666(1994)445
Zukowski, J., see Pawlowska, M. 666(1994)485